The function v(t) is the velocity in m/sec of a particle moving along the x-axis. Determine when the particle is moving to the right, to the left, and stopped.


1)
$$v(t) = 13.6 - 0.2t$$
, $0 \le t \le 120$

2)

Solve the problem.

2) The velocity in m/sec of a particle moving along the x-axis is given by the function $v(t) = 6 \cos 3t$, $0 \le t \le \pi/2$

Find the particle's displacement for the given time interval.



The function v(t) is the velocity in m/sec of a particle moving along the x-axis. Find the total distance traveled by the particle.

3)
$$v(t) = t^2 - 7t + 12, 0 \le t \le 4$$

Solve the problem.

4) A car moving with an initial velocity of 4 mph accelerates at the rate of a(t) = 2.3t mph per

second for 8 seconds. How far did the car travel during those 8 seconds?

It is seconds. How the different travel turning those 8 seconds?

Vel =
$$\int_{0}^{2.3t} 2.3t \, dt \, \frac{mph}{8\pi c}$$

Vel = $\int_{0}^{2.3t} 2.3t \, dt \, \frac{mph}{8\pi c}$
 $\int_{0}^{2.15(64)} - (0) \, dt \, dt \, dt \, dt}$

Vel = $\int_{0}^{2.3t} 2.3t \, dt \, \frac{mph}{8\pi c}$
 $\int_{0}^{2.15(64)} - (0) \, dt \, dt \, dt \, dt}$

Vel = $\int_{0}^{2.15(64)} - (0) \, dt \, dt \, dt}$


Vel = $\int_{0}^{2.15(64)} - (0) \, dt \, dt}$

Vel = $\int_{0}^{2.15(64)} - (0) \, dt \, dt}$

Vel = $\int_{0}^{2.15(64)} - (0) \, dt}$

Vel = $\int_{0}^{$

5) A particle moves along the x-axis (units in cm). Its initial position at t = 0 sec is x(0) = 18. The figure shows the graph of the particle's velocity v(t). The numbers are the areas of the enclosed regions.



6) The rate at which your home consumes electricity is measured in kilowatts. If your home consumes electricity at the rate of 1 kilowatt for 1 hour, you will be charged for 1 "kilowatt-hour" of electricity. Suppose that the average consumption rate for a certain home is modeled by the function $C(t) = 4 - 2.4 \sin(\pi t/12)$, where C(t) is measured in kilowatts and t is the number of hours past midnight. Find the average daily consumption for this home, measured in kilowatt-hours.

7) The following table shows the rate of water flow (in gal/min) from a stream into a pond during a 30-minute period after a thunderstorm. Use the Trapezoidal Rule to estimate the total amount of water flowing into the pond during this period.

Time (min)	Rate (gal/min)	
0	300	- h /
5	350	T= = 1 (y + 2 () + y (x)
10	400	(4.34)
15	350	- /
20	320	T== 2 (-22, 200-2), 2002
25	300	$\frac{1}{2}$
30	250	10(00)/12(00)/1000/
	•	$T = \frac{5}{2} \left(300 + 2(350) + 2(400) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) + 2(350) $
mir	gal	T==== (3990) T== 9975g=1

Find the area of the shaded region.

$$f(x) = -x^{3} + x^{2} + 16x$$

$$g(x) = 4x$$


$$f(x) = -x^{3} + x^{2} + 16x$$

$$f(x) = -x^{3} + x^{3} + x^{2} + 16x$$

$$f(x) = -x^{3} + x^{3} + x^{3} + x^{2} + x^{3} + x^{3}$$


Find the area of the regions enclosed by the lines and curves.


9)
$$y = 129 - 4x^2$$
 and $y = x^2 + 4$



Find the area enclosed by the given curves.

10) Find the area of the region on or above the x-axis bounded by the curves $y^2 = x + 8$ and y = 3x. Find the area by subtracting the area of a triangular region from a larger region.





Find the area of the regions enclosed by the lines and curves.

12) $y^2 = x + 8$ and x = y + 64 $x = y^2 - 8$ y = x - 64 y = x